

octagam® 10%

Immune Globulin
Intravenous (Human) 10%
Liquid Preparation

octagam® 5%

Immune Globulin
Intravenous (Human) 5%
Liquid Preparation

OCTAGAM Co-Pay Program

With the Pfizer OCTAGAM Co-Pay Program, eligible patients may pay as little as **\$0 for OCTAGAM***

- Patients must have commercial insurance to be eligible
- Patients are not eligible if they are enrolled in a state or federally funded insurance program

*Eligible, commercially insured patients may pay as little as \$0 for OCTAGAM and may receive a maximum benefit of \$12,500 per calendar year or the cost of patient's co-pay in a 12-month period (whichever is less) for claims received by the program. Terms and conditions/eligibility requirements apply. See full terms and conditions on page 3 or visit <https://octagam-10.pfizerpro.com/support/octagam-co-payprogram> or <https://octagam-5.pfizerpro.com/support/octagam-co-payprogram>.

See terms and conditions and other details on the following pages.

INDICATIONS AND USAGE FOR OCTAGAM 10%

OCTAGAM 10% is indicated for the treatment of chronic immune thrombocytopenic purpura (cITP) in adults and dermatomyositis (DM) in adults. For patients with cITP, it is used to rapidly increase the platelet count in the blood to help control or prevent bleeding. For patients with DM, it helps improve muscle function and skin rash.

OCTAGAM 10% is a liquid medication that contains Immunoglobulin G (IgG). OCTAGAM 10% is made from human plasma donated by healthy people. OCTAGAM 10% is given through the vein (intravenously) in a hospital, infusion center, or at home.

SELECTED SAFETY INFORMATION FOR OCTAGAM 10%

- Do not use OCTAGAM 10% if you have had a severe allergic reaction to IgG or other blood products or have deficiencies of immunoglobulin A (IgA) with antibodies to IgA.

INDICATION AND USAGE FOR OCTAGAM 5%

OCTAGAM® 5% [Immune Globulin Intravenous (Human)] liquid is indicated for the treatment of primary humoral immunodeficiency (PI).

OCTAGAM 5% is a liquid medication that contains Immunoglobulin G (IgG). OCTAGAM 5% is made from human blood plasma donated by healthy people. OCTAGAM 5% is given through the vein (intravenously) in a hospital, infusion center, or at home.

SELECTED SAFETY INFORMATION FOR OCTAGAM 5%

- Do not use OCTAGAM 5% if you have had a severe allergic reaction to IgG or other blood products or have deficiencies of immunoglobulin A (IgA) with antibodies to IgA.

Please see Selected Safety Information throughout and click [here](#) for Full Prescribing Information, including complete BOXED WARNING, for OCTAGAM 10%. Please see Selected Safety Information throughout and click [here](#) for Full Prescribing Information, including complete BOXED WARNING, for OCTAGAM 5%.



octagam® 10%

Immune Globulin
Intravenous (Human) 10%
Liquid Preparation

octagam® 5%

Immune Globulin
Intravenous (Human) 5%
Liquid Preparation

The Pfizer OCTAGAM Co-Pay Program is available through participating specialty infusion pharmacies



What are OCTAGAM 10% and OCTAGAM 5%?

OCTAGAM 10% and OCTAGAM 5% are liquid medicines for infusion that contain immunoglobulin G (IgG), which is a protein that helps fight infection. OCTAGAM 10% is used to treat adults with chronic immune thrombocytopenic purpura (cITP) and adults with dermatomyositis (DM). OCTAGAM 5% is used to treat primary humoral immunodeficiency (PI).



What are the eligibility requirements for the Pfizer OCTAGAM Co-Pay Program?

- Patients must have commercial insurance to be eligible
- Patients are not eligible if they are enrolled in a state or federally funded insurance program. See full terms and conditions on the next page.



What does the Pfizer OCTAGAM Co-Pay Program provide?

Eligible, commercially insured patients may pay as little as \$0 for OCTAGAM and may receive a maximum benefit of \$12,500 per calendar year or the cost of patient's co-pay in a 12-month period (whichever is less) for claims received by the program. Terms and conditions/eligibility requirements apply. See full terms and conditions on the next page.



Which specialty infusion pharmacies participate in the Pfizer OCTAGAM Co-Pay Program?

All specialty infusion pharmacies may participate in the Pfizer OCTAGAM Co-Pay Program.

READY TO ENROLL?

Contact your specialty infusion pharmacy today.

SELECTED SAFETY INFORMATION FOR OCTAGAM 10%

- OCTAGAM 10% can cause the following:
 - Blood clots in your heart, brain, lungs or other areas of your body
 - Kidney problems, or kidney failure
- Tell your healthcare provider (HCP) if you have an allergy to corn. OCTAGAM 10% contains a type of sugar that is made from corn.
- OCTAGAM 10% can cause the following serious side effects. Contact your HCP if you experience the following:
 - Swelling in your mouth or throat, hives/itching, breathing problems, wheezing, fainting, tightness in your chest, or dizziness. This could be a serious allergic reaction.
 - Decreased urination, swelling in your legs, sudden weight gain, or breathing problems, which could mean kidney failure

SELECTED SAFETY INFORMATION FOR OCTAGAM 5%

- OCTAGAM 5% can cause the following:
 - Blood clots in your heart, brain, lungs, or other areas of your body
 - Kidney problems, or kidney failure
- Do not use OCTAGAM 5% if you are allergic to corn
 - Tell your healthcare provider (HCP) if you have an allergy to corn. OCTAGAM 5% contains a type of sugar that is made from corn
- OCTAGAM 5% can cause the following serious side effects. Contact your HCP if you experience the following:
 - Swelling in your mouth or throat, hives/itching, breathing problems, wheezing, fainting, tightness in your chest, or dizziness. This could be a serious allergic reaction
 - Decreased urination, swelling in your legs, sudden weight gain, or breathing problems, which could be signs of kidney failure

*Please see Selected Safety Information throughout and click [here](#) for Full Prescribing Information, including complete **BOXED WARNING**, for OCTAGAM 10%. Please see Selected Safety Information throughout and click [here](#) for Full Prescribing Information, including complete **BOXED WARNING**, for OCTAGAM 5%.*

octagam® 10%Immune Globulin
Intravenous (Human) 10%
Liquid Preparation**octagam® 5%**Immune Globulin
Intravenous (Human) 5%
Liquid Preparation

TERMS AND CONDITIONS

By using this co-pay card, you acknowledge that you currently meet the eligibility criteria and will comply with the terms and conditions described below:

Patients are not eligible to use this card if they are enrolled in a state or federally funded insurance program, including but not limited to Medicare, Medicaid, TRICARE, Veterans Affairs health care, a state prescription drug assistance program, or the Government Health Insurance Plan available in Puerto Rico (formerly known as "La Reforma de Salud"). Patient must have private insurance. Offer is not valid for cash paying patients. The value of this co-pay card is limited to a maximum of \$12,500 per calendar year or the cost of patient co-pay in a 12-month period, whichever is less. This co-pay card is not valid when the entire cost of your prescription drug is eligible to be reimbursed by your private insurance plan or other private health or pharmacy benefit programs. You must deduct the value of this co-pay card from any reimbursement request submitted to your private insurance plan, either directly by you or on your behalf. You are responsible for reporting use of the co-pay card to any private insurer, health plan, or other third party who pays for or reimburses any part of the prescription filled using the co-pay card, as may be required. You should not use the co-pay card if your insurer or health plan prohibits use of manufacturer co-pay cards. Patient must be 18 years of age or older to be eligible for the co-pay benefit for Octagam 10% and 2 years of age or older to be eligible for the co-benefit for Octagam 5%. This co-pay card is not valid where prohibited by law. The benefit under the co-pay card program is offered to, and intended for the sole benefit of, eligible patients and may not be transferred to or utilized for the benefit of third parties, including, without limitation, third party payers, pharmacy benefit managers, or the agents of either. Co-pay card cannot be combined with any other external savings, free trial or similar offer for the specified prescription (including any program offered by a third party payer or pharmacy benefit manager, or an agent of either, that adjusts patient cost-sharing obligations, through arrangements that may be referred to as "accumulator" or "maximizer" programs). Third party payers, pharmacy benefit managers, or the agents of either, are prohibited from assisting patients with enrolling in the [coupon/co-pay card] program. **Co-pay card will be accepted only at participating pharmacies. If your pharmacy does not participate, you may be able to submit a request for a rebate in connection with this offer. This co-pay card is not health insurance.** Offer good only in the U.S. and Puerto Rico. Co-pay card is limited to 1 per person during this offering period and is not transferable. A co-pay card may not be redeemed more than once per 30 days per patient. No other purchase is necessary. No membership fee. Data related to your redemption of the co-pay card may be collected, analyzed, and shared with Pfizer for market research and other purposes related to assessing Pfizer's programs. Data shared with Pfizer will be aggregated and de-identified; it will be combined with data related to other co-pay card redemptions and will not identify you. Pfizer reserves the right to rescind, revoke or amend this offer without notice. Offer expires 12/31/2024. For more information, call 1-866-642-7606, visit <https://octagam-10.pfizerpro.com/support/octagam-co-payprogram> or write: Octagam Co-Pay Program, P.O. Box 6875, Bridgewater, NJ 08807.

SELECTED SAFETY INFORMATION FOR OCTAGAM 10%

- Pain and/or swelling of an arm or leg with warmth in the affected area, discoloration of an arm or leg, unexplained shortness of breath, chest pain or discomfort that worsens with deep breathing, unexplained rapid pulse, or numbness or weakness on one side of the body; these could be signs of a blood clot.
- Yellow skin or eyes, dark-colored urine, fatigue, or increased heart rate, which could be signs of a blood problem.
- Headache, stiff neck, drowsiness, fever, sensitivity to light, painful eye movements, or nausea and vomiting, which could mean an inflammation of the membranes covering your brain or spinal cord
- Trouble breathing, chest pain, blue lips, arms or legs, and fever, which could be related to a lung problem. This typically occurs 1 to 6 hours following infusion.

Common side effects include headache, fever, nausea, vomiting, increased blood pressure, chills, musculoskeletal pain, dyspnea, infusion site reactions, and increased heart rate.

SELECTED SAFETY INFORMATION FOR OCTAGAM 5%

- Pain and/or swelling of an arm or leg with warmth in the affected area, discoloration of an arm or leg, unexplained shortness of breath, chest pain or discomfort that worsens with deep breathing, unexplained rapid pulse, or numbness or weakness on one side of the body; these could be signs of a blood clot
- Yellow skin or eyes, dark-colored urine, fatigue, or increased heart rate, which could be signs of a blood problem
- Headache, stiff neck, drowsiness, fever, sensitivity to light, painful eye movements, or nausea and vomiting, which could mean an inflammation of the membranes covering your brain or spinal cord
- Trouble breathing, chest pain, blue lips, arms or legs, and fever, which could be related to a lung problem. This typically occurs 1 to 6 hours following infusion

Common side effects include headache and nausea.

Please see Selected Safety Information throughout and click [here](#) for Full Prescribing Information, including complete BOXED WARNING, for OCTAGAM 10%. Please see Selected Safety Information throughout and click [here](#) for Full Prescribing Information, including complete BOXED WARNING, for OCTAGAM 5%.

octagam® 10%

Immune Globulin
Intravenous (Human) 10%
Liquid Preparation

octagam® 5%

Immune Globulin
Intravenous (Human) 5%
Liquid Preparation


OCTAGAM support is available

The following educational resources are available at [OctagamInfo.com](https://www.octagaminfo.com)




My IVIg Therapy Tracker

An easy way for you to record your progress and communicate with your care team



IVIg Infusion Information brochure

Helpful information to prepare you for infusions



What to Expect From Your IVIg Therapy brochure

Helps you prepare for Ig therapy



Ig Companion patient brochure

Information about a free mobile app to help patients on Ig therapy track infusions and side effects

Ig=immunoglobulin;
IVIg=intravenous immunoglobulin.

Your specialty infusion pharmacy may be able to provide you with additional materials about your treatment.

SELECTED SAFETY INFORMATION FOR OCTAGAM 10%

If you use a blood glucose monitor, check with your HCP to ensure that your monitor and test strips are acceptable to use while you are receiving OCTAGAM 10%.

These are not all of the possible side effects with OCTAGAM 10%. Tell your HCPs about any side effects that you have that cause concern or don't go away.

Patients should always ask their doctors for medical advice about adverse events.

You may report an adverse event related to Pfizer products by calling 1-800-438-1985 (US only). If you prefer, you may contact the US Food and Drug Administration (FDA) directly. The FDA has established a reporting service known as MedWatch where healthcare professionals and consumers can report problems they suspect may be associated with the drugs and medical devices they prescribe, dispense, or use. Visit www.fda.gov/MedWatch or call 1-800-FDA-1088.

OCTAGAM® is a registered trademark of Octapharma AG.

Please see Selected Safety Information throughout and click [here](#) for Full Prescribing Information, including complete BOXED WARNING, for OCTAGAM 10%. Please see Selected Safety Information throughout and click [here](#) for Full Prescribing Information, including complete BOXED WARNING, for OCTAGAM 5%.

SELECTED SAFETY INFORMATION FOR OCTAGAM 5%

If you use a blood glucose monitor, check with your HCP to ensure that your type of monitor and test strips can be used while you are receiving OCTAGAM 5%.

These are not all of the possible side effects with OCTAGAM 5%. Tell your HCPs about any side effects that you have.